



DATIM Release 1.29.i was deployed on March 24, 2020.

DATIM 1.29.i includes MER updates

Update:

Updated the language on the TX_PVLS Numerator and Denominator forms for clarity on which disaggregates sum to the total numerator and total denominator.

DSD: TX_PVLS (Numerator)		- Collapse
Auto-Calculate	Number of adults and pediatric patients on ART with suppressed viral load results (<1,000 copies/ml) documented in the medical records and/or supporting laboratory results within the past 12 months. Numerator will auto-calculate from the sum of the Age/Sex/Indication disaggregates.	
Numerator	<input type="text" value="Subtotal"/>	
Required	Disaggregated by Age / Sex / Testing Indication (Fine Disaggregate). Must complete finer disaggregates unless permitted by program.	
Routine		
	Unknown	
Female	Age <1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 50+	<input type="text" value="Subtotal"/>
	Unknown	
Male	Age <1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 50+	<input type="text" value="Subtotal"/>
Routine Subtotal:	<input type="text" value="Subtotal"/>	
Targeted		
	Unknown	
Female	Age <1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 50+	<input type="text" value="Subtotal"/>
	Unknown	
Male	Age <1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 50+	<input type="text" value="Subtotal"/>
Targeted Subtotal:	<input type="text" value="Subtotal"/>	
Conditional	Disaggregated by Age / Sex / Testing Indication (Coarse Disaggregate). USE WITH HQ PERMISSION ONLY [-] Collapse	
	Routine	Targeted
	Unknown Age <15 15+	Unknown Age <15 15+
Female	<input type="text"/>	<input type="text"/>
Male	<input type="text"/>	<input type="text"/>
Sub-total	<input type="text" value="Subtotal"/>	<input type="text" value="Subtotal"/>
Required	Disaggregated by Pregnant/Breastfeeding/Testing Indication. Data on Pregnant/Breastfeeding individuals should be reported in both the "Disaggregated by Pregnant/Breastfeeding/Testing Indication" section and the "Disaggregated by Age/Sex/Testing Indication" section.	
Pregnant	Routine <input type="text"/>	Targeted <input type="text"/>
Breastfeeding	<input type="text"/>	<input type="text"/>
Required	Disaggregated by Key Population/Testing Indication. Data on key populations should be reported in both the "Disaggregated by key population type" section and the "Disaggregated by Age/Sex/Testing Indication" section.	
	Routine	Targeted
PWID	<input type="text"/>	<input type="text"/>
MSM	<input type="text"/>	<input type="text"/>
Transgender People	<input type="text"/>	<input type="text"/>
FSW	<input type="text"/>	<input type="text"/>
People in prison and other closed settings	<input type="text"/>	<input type="text"/>



DSD: TX_PVLS (Denominator) - Collapse

Auto-Calculate Number of adults and pediatric ART patients with a viral load result documented in the medical records and/or supporting laboratory results within the past 12 months. Denominator will auto-calculate from the sum of the Age/Sex/Indication disaggregates.

Denominator Subtotal

Required Disaggregated by Age / Sex / Testing Indication (Fine Disaggregate). Must complete finer disaggregates unless permitted by program.

Routine

Female Unknown Age <1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 50+ Subtotal

Male Unknown Age <1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 50+ Subtotal

Routine Subtotal: Subtotal

Targeted

Female Unknown Age <1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 50+ Subtotal

Male Unknown Age <1 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49 50+ Subtotal

Targeted Subtotal: Subtotal

Conditional Disaggregated by Age / Sex / Testing Indication (Coarse Disaggregate). USE WITH HQ PERMISSION ONLY Collapse

Table with columns for Routine and Targeted, and rows for Female and Male with age groups <15 and 15+.

Required Disaggregated by Pregnant/Breastfeeding/Testing Indication. Data on Pregnant/Breastfeeding individuals should be reported in both the "Disaggregated by Pregnant/Breastfeeding/Testing Indication" section and the "Disaggregated by Age/Sex/Testing Indication" section.

Pregnant Routine Targeted Breastfeeding

Required Disaggregated by Key Population/Testing Indication. Data on key populations should be reported in both the "Disaggregated by key population type" section and the "Disaggregated by Age/Sex/Testing Indication" section.

PWID Routine Targeted MSM Transgender People FSW People in prison and other closed settings